

**Amerifit Nutrition, Inc.**

Estroven .....14

**Beutlich LP**

**Pharmaceuticals**

Peridin-C .....3

**Boston Scientific**

Full Line of Mid Urethral Sling  
Systems (transobturator,  
transvaginal, suprapubic) .....IBC

**Duramed**

**Pharmaceuticals Inc.**

a subsidiary of Barr Laboratories  
Cenestin .....24a-b

**GlaxoSmithKline**

**Consumer Healthcare**

OS-CAL® Calcium Supplement ..9

**Merck & Co., Inc.**

Fosamax Plus D .....OBC

**Pfizer, Inc.**

Estring .....5-6

**Roche Laboratories Inc.  
and GlaxoSmithKline**

Boniva .....16a-d

**Wyeth Pharmaceuticals**

Premarin .....IFC, 1



**ERECTILE DYSFUNCTION**

Sexual problems faced by women during midlife arise from many sources, including a partner's dysfunction. For this reason, the Stat Check in this issue of *Menopause Management* focuses on erectile dysfunction. For more information about the ways in which this disorder affects midlife women, see Clinicians' Forum on page 25.

**15 to 30 million...**

American men and their partners are affected by erectile dysfunction.

*Mayo Clinic, 2004*

**70%...**

of erectile dysfunction cases are related to physical conditions such as diabetes, kidney disease, chronic alcoholism, multiple sclerosis, atherosclerosis, vascular disease, and neurologic disease.

*National Institute of Health, 2003*

**10%...**

or fewer of men who are affected by erectile dysfunction actually receive treatment.

*American Foundation for Urologic Diseases, Inc., 2005*

**35% to 50%...**

of men with diabetes experience erectile dysfunction.

*National Institute of Health, 2003*

**292%...**

is the increase in the number of physician visits made for erectile dysfunction between 1985 and 1999. In 1985 there were 7.7 physician visits for erectile dysfunction per 1,000 men; in 1999 there were 22.5 visits per 1,000 men.

*National Ambulatory Medical Care Survey, 2001*

**3% to 5%...**

of men are estimated to have erectile dysfunction caused by hormone deficiency.

*The Cleveland Clinic Foundation, 2005*

**30%...**

of all cases of erectile dysfunction can be attributed to psychological factors.

*The American Foundation for Urologic Disease Inc., 2005*